

Bayati Eshkaftaki, Ali

Sub-defect of product of $I \times I$ finite sub-defect matrices. (English) [Zbl 1546.15022] Linear Multilinear Algebra 72, No. 10, 1581-1589 (2024).

A square matrix of nonnegative real numbers is said to be *doubly stochastic* if the sum of the elements in each row and column is equal to 1, and *doubly substochastic* if the sum of the elements in each row and column is less than or equal to 1.

L. Cao et al. [Linear Multilinear Algebra 64, No. 11, 2313–2334 (2016; Zbl 1358.15025)] introduced the concept of the sub-defect of an $n \times n$ doubly substochastic matrix A, denoted by sd(A), which is defined as the minimum number of rows and columns needed to be added to A to obtain a doubly stochastic matrix D containing A as a submatrix. Moreover, they showed that the sub-defect of A can be easily calculated using the sum of all elements of A. Later on, L. Cao and S. Koyuncu [Linear Multilinear Algebra 65, No. 4, 653–657 (2017; Zbl 1367.15052)] went on to derive the following lower and upper bounds for the sub-defect of the product of two $n \times n$ doubly substochastic matrices A and B:

 $\max\{\mathrm{sd}(A), \mathrm{sd}(B)\} \leq \mathrm{sd}(AB) \leq \min\{n, \mathrm{sd}(A) + \mathrm{sd}(B)\}.$

In the present article, the author turns his attention on the case of doubly substochastic matrices with infinitely (possibly uncountably) many rows and columns. In this context, the sum of the elements along the rows or columns is to be interpreted as the supremum of the sums over a finite subset of the indexed family of numbers.

In order to briefly describe the main results of the article, a few definitions and explanations are necessary.

- An infinite matrix $[a_{ij}]_{i,j\in\mathcal{I}}$ is called a submatrix of $[b_{ij}]_{i,j\in\mathcal{I}\cup\mathcal{J}}$ if $a_{ij} = b_{ij}$ for all $i, j\in\mathcal{I}$;
- An $\mathcal{I} \times \mathcal{I}$ doubly stochastic matrix D is called a *completion* of an $\mathcal{I} \times \mathcal{I}$ doubly substochastic matrix A if A is a submatrix of D;
- The sub-defect of an infinite doubly substochastic matrix is defined as the minimum cardinal number α such that A has a $(\mathcal{I} \cup \mathcal{J}) \times (\mathcal{I} \cup \mathcal{J})$ completion D, where \mathcal{J} is a set of cardinality α which is disjoint from \mathcal{I} ;
- An infinite matrix A is of finite sub-defect if $sd(A) < \aleph_0$.

The main result achieved by the author is a direct and immediate analogue of that obtained by L. Cao and S. Koyuncu [loc. cit.] for product of doubly substochastic matrices. Indeed, he shows that if A, B are two $\mathcal{I} \times \mathcal{I}$ doubly substochastic matrices of finite sub-defect, then AB is also of finite sub-defect and

 $\max\{\mathrm{sd}(A), \mathrm{sd}(B)\} \leq \mathrm{sd}(AB) \leq \min\{\mathrm{card}(\mathcal{I}), \mathrm{sd}(A) + \mathrm{sd}(B)\}.$

A final remark presents specific examples showing that both bounds are sharp.

Reviewer: Frédéric Morneau-Guérin (Québec)

MSC:

- 15B51 Stochastic matrices
- 15A45 Miscellaneous inequalities involving matrices
- 15A83 Matrix completion problems
- 47B80 Random linear operators
- 03E10 Ordinal and cardinal numbers
- 60E15 Inequalities; stochastic orderings

Keywords:

completion; finite sub-defect; increasable; doubly substochastic; cardinal number

Full Text: DOI

References:

- Bayati Eshkaftaki, A.Increasable doubly substochastic matrices with application to infinite linear equations. Linear Multilinear Algebra. 2022;70(20):5902-5912. doi:. · Zbl 1505.15034
- [3] Bayati Eshkaftaki, A.Row-summable matrices with application to generalization of Schroder's and Abel's functional equations. Aequat Math. 2022;96:525-533. · Zbl 1491.26009
- Fedorov, FM.On the theory of infinite systems of linear algebraic equations. TWMS J Pure Appl Math. 2015;6(2):202-212.
 Zbl 1338.15009
- [5] Hernandez-Pastora, JL.On the solutions of infinite systems of linear equations. Gen Relativ Gravit. 2014;46:1622. doi:. · Zbl 1286.83023
- Shivakumar, PN, Sivakumar, KC, Zhang, Y.Infinite matrices and their recent applications. Switzerland: Springer; 2016. · Zbl 1355.15001
- Cao, L, Koyuncu, S, Parmer, T.A minimal completion of doubly substochastic matrix. Linear Multilinear Algebra. 2016;64(11):2313-2334. · Zbl 1358.15025
- [8] Cao, L.A minimal completion of ((0, 1)-) matrices without total support. Linear Multilinear Algebra. 2018;67:1522-1530. · Zbl 1419.15024
- Cao, L, Chen, Z.Partitions of the polytope of doubly substochastic matrices. Linear Algebra Appl. 2019;563:98-122. · Zbl 1405.15041
- [10] Cao, L, Koyuncu, S.Sub-defect of product of doubly substochastic matrices. Linear Multilinear Algebra. 2017;65(4):653-657.
 · Zbl 1367.15052
- Bahrami, F, Bayati Eshkaftaki, A, Manjegani, SM.Linear preservers of majorization on \(\ell^p(I) \). Linear Algebra Appl. 2012;436:3177-3195.
- [12] Ljubenovic, M, Djordjevic, D.Linear preservers of weak majorization on $((ell^1(I)^+) when I is an infinite set. Linear Algebra Appl. 2017;517:177-198. <math>\cdot$ Zbl 1387.47023

This reference list is based on information provided by the publisher or from digital mathematics libraries. Its items are heuristically matched to zbMATH identifiers and may contain data conversion errors. In some cases that data have been complemented/enhanced by data from zbMATH Open. This attempts to reflect the references listed in the original paper as accurately as possible without claiming completeness or a perfect matching.