

**Radičić, Biljana**

## On geometric circulant matrices with geometric sequence. (English)

Zbl 07895550

**Linear Multilinear Algebra 72, No. 10, 1555-1580 (2024).**

Following an introduction in which he offers a review the literature on  $k$ -circulant matrices, the author presents and motivates the definition of a related class of matrices defined by *C. Kızılateş* and *N. Tuglu* [J. Inequal. Appl. 2016, Paper No. 312, 15 p. (2016; Zbl 1349.15060)], namely the *geometric circulant matrices*, i.e., matrices completely determined by a nonzero complex number  $k \in \mathbb{C} \setminus \{0\}$  and their first row in the sense that they are of the following form:

$$\text{circ}_n\{_{k^*}(c_0, c_1, c_2, \dots, c_{n-1})\} = \begin{pmatrix} c_0 & c_1 & c_2 & \dots & c_{n-2} & c_{n-1} \\ kc_{n-1} & c_0 & c_1 & \dots & c_{n-3} & c_{n-2} \\ k^2 c_{n-2} & kc_{n-1} & c_0 & \dots & c_{n-4} & c_{n-3} \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ k^{n-2} c_2 & k^{n-3} c_3 & k^{n-4} c_4 & \dots & c_0 & c_1 \\ k^{n-1} c_1 & k^{n-2} c_2 & k^{n-3} c_3 & \dots & kc_{n-1} & c_0 \end{pmatrix}.$$

More specifically, the article examines a special case of geometric circulant matrices, where the entries in the first row form a geometric progression of common ratio  $q \in \mathbb{R} \setminus \{0\}$  and of scale factor  $g \in \mathbb{C} \setminus \{0\}$ . In other words, it considers matrices of the form

$$\text{circ}_n\{_{k^*}(g, gq, gq^2, \dots, gq^{n-1})\},$$

and seeks to obtain the determinant, the Frobenius norm, as well as bounds for the spectral norm (i.e., the matrix norm induced by the  $\ell_2$ -norm for vectors) of such matrices.

The main original results set out and demonstrated in this article generally come in pairs:

- An explicit formula (depending on  $k$ ,  $n$  and  $q$ ) is obtained for  $\det(Q)$ , where

$$Q = \text{circ}_n\{_{k^*}(1, q, q^2, \dots, q^{n-1})\}.$$

This formula is then used to obtain an analogous formula for  $\det(G)$ , where

$$G = \text{circ}_n\{_{k^*}(g, gq, gq^2, \dots, gq^{n-1})\}.$$

- The Moore-Penrose inverse of

$$\text{circ}_n\left\{\left(\frac{1}{q^n}\right)^*(1, q, q^2, \dots, q^{n-1})\right\}$$

is obtained for arbitrary natural number  $n > 1$ . This serves as a springboard for the computation of the Moore-Penrose of

$$\text{circ}_n \left\{ \left( \frac{1}{q^n} \right)^* (g, gq, gq^2, \dots, gq^{n-1}) \right\}$$

for arbitrary natural number  $n > 1$ .

- The Moore-Penrose inverse of

$$\text{circ}_n \{_{(-1)^*} (1, -1, 1, \dots, -1, 1) \}$$

is obtained for arbitrary odd natural number  $n > 1$ . Once again, this serves as a springboard for the computation of the Moore-Penrose of

$$\text{circ}_n \{_{(-1)^*} (g, -g, g, \dots, -g, g) \}$$

for arbitrary odd natural number  $n > 1$ .

Lastly, and perhaps most interestingly, explicit formulas for the Frobenius norm of  $\text{circ}_n \{_{k^*} (g, gq, gq^2, \dots, gq^{n-1}) \}$  are derived, along with upper and lower bounds for their spectral norm.

It is worth highlighting that the article contains a number of numerical examples to aid understanding.

Reviewer: Frédéric Morneau-Guérin (Québec)

#### MSC:

- 15B05 Toeplitz, Cauchy, and related matrices
- 15A09 Theory of matrix inversion and generalized inverses
- 15A15 Determinants, permanents, traces, other special matrix functions
- 15A60 Norms of matrices, numerical range, applications of functional analysis to matrix theory

#### Keywords:

geometric circulant matrix; geometric sequence; determinant of a matrix; norms of a matrix; generalized inverses of a matrix

[PDF](#) [BibTeX](#) [XML](#) [Cite](#)

Full Text: [DOI](#)

#### References:

- [1] Shen, SQ, Cen, JM. On the bounds for the norms of r-circulant matrices with the Fibonacci and Lucas numbers. *Appl Math Comput.* 2010;216:2891-2897. · Zbl 1211.15029
- [2] He, C, Ma, J, Zhang, K, et al. The upper bound estimation on the spectral norm of r-circulant matrices with the Fibonacci and Lucas numbers. *J Inequal Appl.* 2015;2015:72. · Zbl 1314.15024
- [3] Köme, C, Yazlik, Y. On the spectral norms of r-circulant matrices with the biperiodic Fibonacci and Lucas numbers. *J Inequal Appl.* 2017;2017:192. · Zbl 1372.15013
- [4] Köme, C, Yazlik, Y. On the determinants and inverses of r-circulant matrices with the biperiodic Fibonacci and Lucas numbers. *Filomat.* 2018;32(10):3637-3650. · Zbl 1499.15103

- [5] Radičić, B.On k-circulant matrices with arithmetic sequence. *Filomat.* 2017;31(8):2517-2525. · Zbl 1488.15056
- [6] Radičić, B.On k-circulant matrices (with geometric sequence). *Quaest Math.* 2016;39(1):135-144. · Zbl 1422.15012
- [7] Radičić, B.On k-circulant matrices involving geometric sequence. *Hacet J Math Stat.* 2019;48(3):805-817. · Zbl 1488.15055
- [8] Kizilates, C, Tuglu, N.On the bounds for the spectral norms of geometric circulant matrices. *J Inequal Appl.* 2016;2016:312. · Zbl 1349.15060
- [9] Böttcher, A, Silbermann, B.*Introduction to large truncated toeplitz matrices.* Springer; 1999. · Zbl 0916.15012
- [10] Garoni, C, Serra-Capizzano, S.*Generalized locally toeplitz sequences: theory and applications – volume I.* Springer; 2017. · Zbl 1376.15002
- [11] Gray, RM.*Toeplitz and circulant matrices: A review.* *Found Trends Commun Inf Theory.* 2006;2(3):155-239.
- [12] Iohvidov, IS.*Hankel and Toeplitz matrices and forms: algebraic theory.* Boston: Birkhäuser; 1982. · Zbl 0493.15018
- [13] Ng, MK.*Iterative methods for toeplitz systems.* Oxford University Press; 2004. · Zbl 1059.65031
- [14] Shi, B.The spectral norms of geometric circulant matrices with the generalized k-Horadam numbers. *J Inequal Appl.* 2018;2018:14. · Zbl 1381.15011
- [15] Uygun, S, Aytar, H.On the bounds for the spectral norms of geometric and r-circulant matrices with bi-periodic Jacobsthal numbers. *J Appl Math Inform.* 2020;38(1-2):99-112. · Zbl 1463.15047
- [16] Zielke, G.Some remarks on matrix norms, condition numbers, and error estimates for linear equations. *Linear Algebra Appl.* 1988;110:29-41. · Zbl 0654.15018
- [17] Penrose, R.A generalized inverse for matrices. *Proc Cambridge Philos Soc.* 1955;51:406-413. · Zbl 0065.24603
- [18] Ben-Israel, A, Greville, TNE.*Generalized inverses: theory and applications.* New York: John Wiley and Sons, Springer; 1974. · Zbl 0305.15001
- [19] Piziak, R, Odell, PL.*Full rank factorization of matrices.* *Math Mag.* 1999;72(3):193-201. · Zbl 1006.15009
- [20] Puntanen, S, Styan, GPH, Isotalo, J.*Matrix tricks for linear statistical models.* Berlin: Springer; 2011. · Zbl 1291.62014
- [21] Radičić, B.Circulant matrices with a special (geometric and arithmetic) sequence. *Indian J Math.* 2016;58(1):1-16. · Zbl 1343.15018
- [22] Bueno, ACF.Right circulant matrices with geometric progression. *Int J Appl Math Res.* 2012;1(4):593-603.
- [23] Horn, RA, Johnson, CR.*Topics in matrix analysis.* Cambridge: Cambridge Univ. Press; 1991. · Zbl 0729.15001
- [24] Horn, RA.The Hadamard product. *Proc Sympos Appl Math.* 1990;40:87-169. · Zbl 0712.15011
- [25] Liu, S, Trenkler, G.Hadamard, Khatri-Rao, Kronecker and other matrix products. *Int J Inf Syst Sci.* 2008;4(1):160-177. · Zbl 1159.15008

This reference list is based on information provided by the publisher or from digital mathematics libraries. Its items are heuristically matched to zbMATH identifiers and may contain data conversion errors. In some cases that data have been complemented/enhanced by data from zbMATH Open. This attempts to reflect the references listed in the original paper as accurately as possible without claiming completeness or a perfect matching.